

Figure 4. Plot showing locations of Mars Global Surveyor (MGS) (>3 m/pixel) and Mars Reconnaissance Orbiter (HiRISE) (25 cm/pixel) images in the map region obtained during all seasons and released as of May 2009, on a background showing the topography in shaded relief based on the MOLA DEM.




Figure 5. Composite mosaic of three color images acquired by the Mars Reconnaissance Orbiter Mars Color Imager (MARCI) on October 16th of 2006 during Martian northern summer, solar longitude, $L_s = 121^\circ$. The MARCI frames are blended with a MOGA shaded relief (-250 m/pix) to enhance topographic features, which are sometimes muted by atmospheric haze in the color images. The image shows the region north of lat 72° N, at -1000 pixels. The color is derived from three spectral bands acquired at 425 nm, 550 nm, and 600 nm. We mapped the high-albedo residual ice using this mosaic. Additional color units shown in Tanaka and others (2008) include beige-colored dust, low-albedo basaltic sand, and mixtures of low- and moderate-albedo materials with sparse residual ice.




Figure 6. Map of unconformities (red lines) and grabens (black lines with balls in box) in *Platum Boreum* 1 unit and *Platum Boreum* 3. Background shows the topography in shaded relief based on the MOLA DEM as well as unit contacts.




Figure 7. Plot showing the distribution of impact craters > 2 km in diameter in the map region (see table 4 regarding morphologic and other characteristics). These include craters used in crater statistics (table 2 and fig. 8).




Figure 8 consists of two panels. The top panel is a log-log plot of Crater count (km⁻²) versus Diameter (km). It shows three data series: Plutonium-Boreum rapids unit (red), Borealis impact unit (green), and media region unit (blue). The bottom panel shows several plots for the same units: R-squared density, C, incremental density, and log in square root of the diameter increments, D₂, production-function fit, Cumulative, and areal plots. The plots are color-coded to match the units in the top panel.

Geologic Map of the North Polar Region of Mars

By
Kenneth L. Tanaka and Corey M. Fortezzo
2012


Prepared on behalf of the Planetary Geology and Geophysics Program
Solar System Exploration Division, Office of Space Science, National
Aeronautics and Space Administration
Edited by Theresa Iki
Cartography by Darlene A. Ryan
Manuscript approved for publication August 29, 2011

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

For sale by U.S. Geological Survey, Information Service, Box 25208, Federal Center, Denver, CO 80225. 1-888-ASK-1935

Digital files available at <http://pubs.usgs.gov/pdfiles/3737/>

Suggested citation: TANAKA, K.L., and FORTNEY, C.M., 2013. Geologic map of the North Polar Region of Mars: The U.S. Geological Survey Scientific Investigations Map 3117, scale 1:2,500,000 (<http://pubs.usgs.gov/pdfiles/3737/>).

 PRINTED on recycled paper




Figure 3 (right). Mars Odyssey (ODY) mission Thermal Emission Imaging System (THEMIS) visual-range image mosaics of the Planum Boreum region in Polar Stereographic projection (courtesy of P.R. Christensen, Arizona State University). The mosaics are at half the spatial resolution of the original images, which were obtained at full (18 m/pixel) and half instrument resolution. *A*, Summertime images at 36 m/pixel north of lat 86° N. *B*, Summertime images at 72 m/pixel north of lat 75° N. *C*, Springtime images at 36 m/pixel north of lat 75° N. *D*, Springtime images at 72 m/pixel north of lat 75° N.